

A set of invariant quality factors measuring the deviation from the Kerr metric

Alfonso García-Parrado Gómez-Lobo ^{*}
 Centro de Matemática, Universidade do Minho
 4710-057 Braga, Portugal

José M. M. Senovilla [†]
 Física Teórica, Universidad del País Vasco
 Apartado 644, E-48080 Bilbao, Spain

November 30, 2012

Abstract

A number of scalar invariant characterizations of the Kerr solution are presented. These characterizations come in the form of *quality factors* defined in stationary space-times. A quality factor is a scalar quantity varying in the interval $[0, 1]$ with the value 1 being attained if and only if the space-time is locally isometric to the Kerr solution. No knowledge of the Kerr solution is required to compute these quality factors. A number of different possibilities arise depending on whether the space-time is Ricci-flat and asymptotically flat, just Ricci-flat, or Ricci non-flat. In each situation a number of quality factors are constructed and analysed. The relevance of these quality factors is clear in any situation where one seeks a rigorous formulation of the statement that a space-time is “close” to the Kerr solution, such as: its non-linear stability problem, the asymptotic settlement of a radiating isolated system undergoing gravitational collapse, or in the formulation of some uniqueness results.

1 Introduction

The Kerr metric [16] is a fundamental exact solution in General Relativity providing the unique set of models describing the region outside a rotating black hole, so that its physical relevance is unquestionable. The uniqueness properties of the Kerr solution stem from powerful mathematical results stating, roughly speaking, that it is the only analytic, asymptotically flat vacuum solution of the Einstein field equations (without cosmological constant) representing a black hole with a *regular* event horizon (a precise, up-to-date formulation of this uniqueness result and its proof can be found in [5, 6]). The generalization of this result for non-analytic solutions of the field equations has been an open question for many years and recent progress towards its solution has been reported in [14, 1, 15]

A completely open problem regarding the Kerr solution is its *stability* under *nonlinear* perturbations. The non-linear stability of the Kerr solution —to be more precise, the non-linear stability of its *domain of outer communication*— can be formulated as the global existence of a solution of the vacuum Einstein equations corresponding to initial data whose maximal Cauchy development is *close* to a subset of the Kerr’s domain of outer communication. Results in this direction already exist for other well known exact solutions to the Einstein field equations [8, 9, 10, 4, 17, 20] and therefore, given the relevance of the Kerr solution, it is important attempting to obtain similar results for it. In this regard, it is crucial to establish what is the precise meaning of the statement that a solution of the field equations be “close to the Kerr solution”.

^{*}E-mail address: alfonso@math.uminho.pt

[†]E-mail address: josemm.senovilla@ehu.es

Another point of relevance of the Kerr solution arises from its status as the stationary equilibrium limit reached in the evolution of an isolated system undergoing gravitational collapse. Due to its uniqueness properties just mentioned, it is widely believed that after the gravitational wave content has been radiated away such systems will eventually settled down as the exterior region of a Kerr solution. Any rigorous proof of this statement requires to introduce the notion of “approaching the (exterior) of the Kerr solution” which in turn relies on a further notion of “being close to Kerr”. With the aid of this notion one could devise numerical simulations of a radiating isolated system and test whether in the asymptotic regime it is close to the Kerr solution.

The purpose of the present work is to put forward a number of *local and invariant* criteria which enable us to answer the question of when a given space-time is “close” to the Kerr solution. By local and invariant we mean that the criteria are formulated in terms of scalar quantities, so that they have a perfectly defined value at each point of the space-time, and are independent of the coordinate systems used. This makes them suitable for their application to the problems discussed in the previous paragraphs. Our approach consists in introducing *quality factors* to measure the deviation of a given solution from the Kerr space-time. A quality factor is a dimensionless scalar quantity varying in the interval $[0, 1]$ and such that the unity value is attained on a neighborhood of a point if and only if the space-time is isometric to the Kerr solution there. Ideally one seeks quality factors which are constructed from invariantly defined quantities of the geometry. The existence of such type of quality factors permits to quantify invariantly, by a normalized non-negative real number, “how close” to the Kerr solution is the space-time under consideration.

These quality factors are obtained from a number of coordinate-free characterisations of the Kerr solutions. A coordinate-free characterisation of the Kerr solution is a set of geometric conditions on a space-time which are fulfilled if and only if the space-time is locally isometric to the Kerr solution. A number of them have been presented in the literature, and we rely specifically on those which are formulated as local conditions, computable at any given point. An example of these can be found in [7] where the Kerr metric is characterised in terms of a set of conditions on the concomitants of the curvature tensor at a point, so they are fully invariant. Another local invariant characterisation of the Kerr solution was put forward in [21]. In this case one assumes the existence of a Killing vector in the space-time (an invariant property) and that the space-time is a vacuum solution. The characterization is then written as a set of local conditions involving the Killing vector and the curvature tensor. This result is the starting point of our work. The plan is to cast this invariant characterisation as a single scalar condition having the properties of a quality factor. Indeed, it turns out that this can be done in more than one way and we present a number of quality factors which are relevant in different situations.

An important property of our approach is that the quality factors that we introduce depend exclusively on invariant geometric properties of the space-time under study. One does not need to know anything about the Kerr solution, nor one has to put the space-time in relation with the Kerr solution (by diffeomorphisms, for instance). Actually, the quality factors can be explicitly computed for any given stationary space-time.

The question of whether or not our quality factors are good enough to measure the deviation of solutions from the Kerr metric will have to be evaluated, and a calibration of the meaning of any quality factor having values close to one (say 0.95 or more) should be carefully and exhaustively performed. In this sense, we support our analysis with the study of some practical examples where the quality factors have been explicitly computed. The results are in agreement with intuitive expectations of when a space-time can be considered closed to the Kerr solution.

The plan of this work is as follows: in section 2 we recall some generalities of vacuum space-times possessing a Killing vector. In sections 3 and 4 we construct a quality factor valid for any stationary, Ricci-flat and asymptotically flat space-time. This quality factor is based on the so-called *Mars-Simon* tensor. The more general case of stationary space-times is addressed in section 5 where we construct a number of quality factors based on the spacetime *Simon tensor*. We discuss quality factors valid for vacuum solutions and a quality factor which applies to non-vacuum solutions (although in this case the quality factor being unity is just a necessary condition for the space-time to be locally isometric to the Kerr solution). In section 6, we carried out numerical computations of the quality factors in practical cases with the aid of the system *xAct* [26, 27]. Finally we discuss some open issues of our analysis in section 7.

2 Geometric preliminaries

Let $\mathcal{M}$ be a 4-dimensional smooth manifold endowed with a smooth Lorentzian metric g_{ab} (signature convention $(-, +, +, +)$). The pair $(\mathcal{M}, g_{ab})$ will be referred to as a *Lorentzian manifold* or *spacetime*. Small Latin letters $a, b, c, \dots$ will be used to denote abstract indices for tensors fields on any tensor bundle built from $T(\mathcal{M})$ and its dual $T^*(\mathcal{M})$. Square (rounded) brackets enclosing a group of indices will denote the antisymmetrization (symmetrization) operation. The inverse of g_{ab} is g^{ab} and these quantities can be used to raise and lower tensor indices in the standard way. The Levi-Civita covariant derivative compatible with g_{ab} is the operator ∇_a , while $R_{abc}{}^d$ and R_{ab} denote respectively the Riemann and Ricci tensors arising from ∇_a . Our conventions in the definitions of these quantities are set by the relations,

$$\nabla_a \nabla_b Z^c - \nabla_b \nabla_a Z^c = R_{bad}{}^c Z^d, \quad R_{ab} \equiv R_{acb}{}^c,$$

where the first relation is just the Ricci identity valid for any vector field Z^c . Sometimes we will denote tensorial quantities with index-free notation in which case boldface capital letters will be used (the rank of the tensorial quantity should then be clear from the context). The complex conjugate of a complex quantity will be denoted with an overbar.

A vector field $\vec{\zeta}$ on $\mathcal{M}$ is called a *Killing vector* if it fulfills the differential condition

$$\mathcal{L}_{\vec{\zeta}} g_{ab} = 0. \quad (1)$$

where $\mathcal{L}_{\vec{\zeta}}$ denotes the Lie derivative with respect to $\vec{\zeta}$. The previous expression can be written in terms of the Killing 1-form $\zeta_a \equiv g_{ab} \zeta^b$ in the well-known equivalent form

$$\nabla_a \zeta_b + \nabla_b \zeta_a = 0. \quad (2)$$

From now on we will assume that the spacetime on which we work possesses a Killing vector. Killing vectors have been extensively studied and their properties can be found in many places of the literature but in order to make our presentation as self-contained as possible we will review here those properties needed in our work. First of all we assume that the Killing vector $\vec{\zeta}$ is a smooth vector field in $\mathcal{M}$ (indeed, the smoothness of the Killing vector follows if one just assumes that it is a C^2 vector field, see below). The condition (2) implies that ζ_a has the property

$$\nabla_{[a} \zeta_{b]} = \nabla_a \zeta_b,$$

which enables us to define an exact 2-form, called the Killing 2-form, by $F_{ab} \equiv \nabla_a \zeta_b$. It follows immediately that

$$\nabla_{[a} F_{bc]} = 0. \quad (3)$$

Using the Killing condition (2) and the Ricci identity one may compute the covariant derivative of the Killing 2-form, yielding

$$\nabla_b F_{ac} = -R_{acbd} \zeta^d. \quad (4)$$

The norm of the Killing vector is $\zeta_a \zeta^a$ and no restriction on it is imposed at this stage. If we set $\lambda \equiv \zeta_a \zeta^a$ then the Killing condition (2) easily gives

$$\nabla_b \lambda = 2\zeta^a \nabla_b \zeta_a = 2F_{ba} \zeta^a. \quad (5)$$

The identities discussed so far hold regardless of the space-time dimension and its matter content. In the particular case of Ricci-flat 4-dimensional space-time we have additional properties to be detailed next. First of all, in a Ricci-flat space-time, that is with vanishing Ricci tensor $R_{ab} = 0$, the Riemann tensor coincides with the Weyl tensor C_{abcd} which in four dimensions has the algebraic properties $C_{abcd}^* = {}^*C_{abcd}$ where the Weyl tensor right and left duals are defined respectively by

$$C_{abcd}^* \equiv \frac{1}{2} \eta_{cdpq} C_{ab}{}^{pq}, \quad {}^*C_{abcd} \equiv \frac{1}{2} \eta_{abpq} C_{cd}{}^{pq},$$

and η_{abcd} is the canonical volume 4-form. Therefore, we will only use the Weyl tensor and its right dual from now on. Also, in four dimensions one can define the dual of the Killing 2-form F_{ab}

by $F_{ab}^* \equiv F^{pq}\eta_{abpq}/2$. From F_{ab}^* we introduce the twist 1-form $\omega_a \equiv F_{ab}^*\zeta^b$ which, for a Ricci-flat spacetime, can be shown to be a closed 1-form, $\nabla_{[a}\omega_{b]} = 0$. The *local* potential arising from ω_a is called the *twist potential* ω . The 2-form F_{ab}^* has the differential properties

$$\nabla_c F_{ab}^* = C_{abcd}^* \zeta^d \Rightarrow \nabla_{[a} F_{bc]}^* = 0. \quad (6)$$

It is possible to achieve a great simplification in many computations by introducing *complex quantities*. In this way we define the *self-dual* Weyl tensor, and the self-dual Killing 2-form by the expressions

$$\mathcal{C}_{abcd} \equiv C_{abcd} + i C_{abcd}^*, \quad \mathcal{F}_{ab} \equiv F_{ab} + i F_{ab}^*. \quad (7)$$

These quantities fulfill the well-known algebraic properties

$$\mathcal{C}_{abpq}^* \equiv \frac{1}{2}\eta_{pqcd}\mathcal{C}_{ab}{}^{cd} = -i\mathcal{C}_{abpq}, \quad \mathcal{F}_{ab}^* \equiv \frac{1}{2}\eta_{abcd}\mathcal{F}^{cd} = -i\mathcal{F}_{ab} \quad (8)$$

and the identities

$$\mathcal{F}_{ac}\mathcal{F}_b{}^c = \frac{1}{4}(\mathcal{F} \cdot \mathcal{F})g_{ab}, \quad \mathcal{F}_{ac}\overline{\mathcal{F}}_b{}^c = \mathcal{F}_{bc}\overline{\mathcal{F}}_a{}^c, \quad \mathcal{F} \cdot \mathcal{F} \equiv \mathcal{F}_{ab}\mathcal{F}^{ab}. \quad (9)$$

With the aid of these new quantities, one can write the differential conditions fulfilled by F_{ab} and F_{ab}^* in a compact form

$$\nabla_c \mathcal{F}_{ab} = -\mathcal{C}_{abcd}\zeta^d, \quad \nabla_{[c}\mathcal{F}_{ab]} = 0. \quad (10)$$

Also, if we introduce the Ernst 1-form

$$\sigma_a \equiv 2\mathcal{F}_{ab}\zeta^b, \quad (\sigma_a\zeta^a = 0)$$

one has that in a Ricci-flat space-time

$$\nabla_{[a}\sigma_{b]} = 0, \quad (11)$$

which means that one can find a local potential σ for σ_a , called the Ernst potential, which can be defined in terms of the Killing norm and twist as

$$\sigma \equiv 1 + \lambda + 2i\omega \Rightarrow \nabla_a\sigma = \sigma_a. \quad (12)$$

More generally, the Ernst 1-form is known to be closed in any space-time that satisfies $\zeta_{[c}R_{a]b}\zeta^b = 0$. The reader is warned about the non-standard choice of the real part of σ in (12), as this is universally chosen to be λ in the literature. However, our simple choice will simplify the appearance of many expressions below. Observe that σ is defined up to the addition of an additive purely imaginary constant, so that we have the gauge freedom

$$\sigma \longrightarrow \sigma' = \sigma + i\alpha, \quad \alpha \in \mathbb{R}. \quad (13)$$

This freedom will be relevant later when we deal with spacetimes locally, without reference to their possible asymptotic properties.

The quantities $\mathcal{F}_{ab}$ and σ_a can be related by the formula

$$-\lambda\mathcal{F}_{ab} = \zeta_{[a}\sigma_{b]} + \frac{i}{2}\eta_{abcd}\zeta^c\sigma^d. \quad (14)$$

See [13, 23] for a more complete review of the properties of the quantities just introduced.

3 The Mars-Simon tensors and their properties

At those points of $\mathcal{M}$ in which $\sigma \neq 0$ we can introduce the following rank-4 tensor

$$\mathcal{S}_{abcd} \equiv \mathcal{C}_{abcd} + \frac{6}{\sigma} \left(\mathcal{F}_{ab}\mathcal{F}_{cd} - \frac{\mathcal{I}_{abcd}}{3}\mathcal{F} \cdot \mathcal{F} \right), \quad (15)$$

where

$$\mathcal{I}^{ab}_{cd} \equiv \frac{1}{4}(\mathbf{i} \eta^{ab}_{cd} + \delta^a_c \delta^b_d - \delta^a_d \delta^b_c) \quad (16)$$

is the identity tensor in the space of self-dual 2-forms. Observe that $\mathcal{S}_{abcd}$ is affected by the freedom (13) so that one has to either (i) try to choose a preferred Ernst potential, or (ii) add the arbitrary constant α and consider one-parameter families of potentials $\sigma(\alpha)$ and of tensors $\mathcal{S}_{abcd}(\alpha)$.

The tensor $\mathcal{S}_{abcd}$ was first introduced in [14]¹ and it was called the *Mars-Simon tensor*, a terminology which we also adopt here. This tensor satisfies a number of remarkable algebraic and differential identities whose proof can be found in [14, 11]. The most relevant ones for us come from the fact that the Mars-Simon tensor is a *Weyl field* (also called a Weyl candidate), which means that it fulfills the same algebraic properties as the Weyl tensor

$$\mathcal{S}_{abcd} = -\mathcal{S}_{bacd}, \quad \mathcal{S}_{abcd} = \mathcal{S}_{cdab}, \quad \mathcal{S}_{abcd} + \mathcal{S}_{acdb} + \mathcal{S}_{adbc} = 0, \quad \mathcal{S}^a_{bac} = 0 \quad (17)$$

and it is self-dual

$$\mathbf{i} \mathcal{S}^*_{abcd} = \mathbf{i}^* \mathcal{S}_{abcd} = \mathcal{S}_{abcd}. \quad (18)$$

Its covariant divergence is given in Ricci-flat spacetimes by [14, 11]

$$\sigma \nabla_a \mathcal{S}^a_{bcd} = 2\zeta^a (2(\mathcal{F}^p_{[d} \mathcal{S}_{c]bap} + \mathcal{F}_b^p \mathcal{S}_{dcap}) + g_{b[d} \mathcal{S}_{c]apq} \mathcal{F}^{pq}). \quad (19)$$

We note the important property that this covariant divergence is a linear expression in the Mars-Simon tensor itself. Using this one can further show that $\mathcal{S}_{abcd}$ fulfills a hyperbolic equation [14] and that it vanishes in a globally hyperbolic space-time if it is zero on a Cauchy hypersurface (causal propagation) [11].

The tensor $\mathcal{S}_{abcd}$ plays an important role in characterizing invariantly the Kerr solution. One of such characterizations requires that the space-time be asymptotically flat as originally formulated by Mars in [21], Theorem 1 below. To that purpose, we briefly introduce the concept of stationary asymptotically flat end.

3.1 Asymptotically flat end $\mathcal{M}_\infty$

Assume that there exists a submanifold $\mathcal{M}_\infty \subset \mathcal{M}$ diffeomorphic to $\mathbb{R} \times \{x \in \mathbb{R}^3 : |x| > R\}$ for some R large enough. Suppose further, that in the local Cartesian coordinates (t, x^i) , $i = 1, 2, 3$ on $\mathcal{M}_\infty$ defined by the diffeomorphism we have $\vec{\zeta} = \partial_t$ (time translation at infinity) and also the conditions

$$|g_{ab} - \eta_{ab}| + |r \partial_{x^i}(g_{ab})| \leq A r^{-B}, \quad \partial_t g_{ab} = 0, \quad (20)$$

where A and B are positive constants, g_{ab} and η_{ab} represent, respectively, the components of the metric and the Minkowski metric in the above local coordinates and $r \equiv \sqrt{(x^1)^2 + (x^2)^2 + (x^3)^2}$. A set $\mathcal{M}_\infty$ with these properties is called a *stationary asymptotically flat end*. It is well-known (see e.g. [2]) that, using the vacuum Einstein equations (without cosmological constant), the components of the metric tensor have the following asymptotic behaviour in $\mathcal{M}_\infty$

$$g_{tt} = -1 + \frac{2M}{r} + O(r^{-2}), \quad g_{tx^i} = -\epsilon_{ijk} \frac{4S^i x^k}{r^3} + O(r^{-3}), \quad g_{x^i x^j} = \delta_{ij} + O(r^{-1}), \quad (21)$$

where M is the Komar mass [18] of the Killing vector $\vec{\zeta}$ in $\mathcal{M}_\infty$, ϵ_{ijk} is the Levi-Civita symbol in dimension 3, and δ_{ij} is the Kronecker symbol (Riemannian flat metric) in dimension 3.

3.2 Two theorems on $\mathcal{S}_{abcd}$

We are now ready to present the first Mars' theorem.

¹The definition of [14] uses a different choice of the Ernst potential.

Theorem 1. *Let $(\mathcal{M}, g_{ab})$ be a smooth Ricci-flat spacetime possessing a Killing vector $\vec{\zeta}$ and a stationary asymptotically flat end $\mathcal{M}_\infty$ such that $\vec{\zeta}$ tends to a time translation at infinity and the Komar mass of $\vec{\zeta}$ at $\mathcal{M}_\infty$ does not vanish. Fix the freedom (13) by letting $\sigma \rightarrow 0$ at $\mathcal{M}_\infty$ and construct the tensor $\mathcal{S}_{abcd}$ according to (15).*

If $\mathcal{S}_{abcd}$ vanishes, then $(\mathcal{M}, g_{ab})$ is locally isometric to the Kerr spacetime.

This theorem includes a global condition, namely, the existence of a stationary asymptotically flat end $\mathcal{M}_\infty$. This is important to fix the freedom (13) for the potential σ to be used in the definition of $\mathcal{S}_{abcd}$, as one can choose the potential such that σ goes to 0 at the asymptotically flat end fixing the constant α in (13) so that Theorem 1 possesses a totally unambiguous meaning.

Nevertheless, it is possible to formulate a purely local characterization of the Kerr metric, see theorem 1 in [22]. To that end, we need to control the change of $\mathcal{S}_{abcd}$ under (13), and thus we introduce the tensor

$$\mathcal{Q}_{abcd} \equiv 6 \left(\mathcal{F}_{ab} \mathcal{F}_{cd} - \frac{\mathcal{I}_{abcd}}{3} \mathcal{F} \cdot \mathcal{F} \right). \quad (22)$$

The algebraic properties of $\mathcal{F}_{ab}$ and $\mathcal{I}_{abcd}$ guarantee that $\mathcal{Q}_{abcd}$ is a Weyl candidate too. It is a very simple matter to check that under the freedom (13) the Mars-Simon tensor changes as

$$\mathcal{S}'_{abcd} = \mathcal{S}_{abcd} - \frac{i\alpha}{\sigma(\sigma + i\alpha)} \mathcal{Q}_{abcd} \quad (23)$$

Therefore the existence of a potential such that the Mars-Simon tensor vanishes is equivalent to the proportionality of $\mathcal{S}_{abcd}$ —constructed from *any* given $\sigma \neq 0$ — and $\mathcal{Q}_{abcd}$, with some given properties of the proportionality function. This will be relevant in section 5 and allowed Mars to prove his second, purely local, theorem.

Theorem 2. *Let $(\mathcal{M}, g_{ab})$ be a smooth non-flat Ricci-flat spacetime possessing a Killing vector $\vec{\zeta}$ and assume further that the tensor $\mathcal{S}_{abcd}$ is well defined on $\mathcal{M}$. If there exists a potential σ' in (13) such that the following conditions are fulfilled on $\mathcal{M}$*

$$\mathcal{S}'_{abcd} = 0, \quad \mathcal{F} \cdot \mathcal{F} = -\frac{\sigma'^4}{4M^2}, \quad M \in \mathbb{R} \setminus \{0\}, \quad (24)$$

then $\mathcal{M}$ is locally isometric to the Kerr space-time.

Remark 1. The constant M here is the Komar mass of $\vec{\zeta}$ as introduced in subsection 3.1. Note that the imposed condition $M \neq 0$ on the Komar mass excludes the flat Minkowski solution even though strictly speaking it belongs to the Kerr family. In this work “Kerr space-time” will mean any member of the Kerr family with non-zero mass parameter.

3.3 The space-time Simon tensor

The vanishing of $\mathcal{S}'_{abcd}$ is of course equivalent to a particular proportionality between $\mathcal{C}_{abcd}$ and $\mathcal{Q}_{abcd}$:

$$\mathcal{S}'_{abcd} = 0 \quad \Longleftrightarrow \quad \mathcal{C}_{abcd} = -\frac{1}{\sigma'} \mathcal{Q}_{abcd} \quad \Longleftrightarrow \quad \exists \alpha \in \mathbb{R} : \mathcal{C}_{abcd} = -\frac{1}{\sigma + i\alpha} \mathcal{Q}_{abcd} \quad (25)$$

and this is why it will be convenient, and very useful, to use another tensor which measures such a proportionality. To that end, let us define

$$\mathcal{S}_{abc} \equiv \gamma_{a[b} \mathcal{C}_{c]mrd} \zeta^m \mathcal{F}^{rd} + 4\zeta^m \zeta^r \mathcal{C}_{mar[c} \sigma_{b]} \quad (26)$$

where here and in the rest of the paper we use the abbreviation

$$\gamma_{ac} \equiv \zeta_a \zeta_c - \lambda g_{ac}. \quad (27)$$

The tensor $\mathcal{S}_{abc}$ is called the *space-time Simon tensor* [21] and it can be defined in any spacetime with a Killing vector, independently of the existence of a potential σ for σ_a . This will be very important later. $\mathcal{S}_{abc}$ has the algebraic properties of a Lanczos potential

$$\mathcal{S}^b_{bc} = 0, \quad \mathcal{S}_{a[bc]} = \mathcal{S}_{abc}, \quad \mathcal{S}_{abc} + \mathcal{S}_{bca} + \mathcal{S}_{cab} = 0 \quad (28)$$

hence the space-time Simon tensor is a *Lanczos field* (or a Lanczos candidate). From (26) we deduce the additional properties

$$\zeta^a \mathcal{S}_{abc} = 0, \quad \zeta^b \mathcal{S}_{abc} = 0 \quad (29)$$

so that $\mathcal{S}_{abc}$ is totally *orthogonal* to the Killing vector $\vec{\zeta}$ from which it is constructed.

A fundamental property of this tensor is (see considerations leading to eq. (16) in [21]).

Lemma 1. *In arbitrary spacetimes with a non-null Killing vector $\vec{\zeta}$ such that $\mathcal{F} \cdot \mathcal{F} \neq 0$, the vanishing of the space-time Simon tensor is equivalent to the proportionality of the self-dual tensors $\mathcal{C}_{abcd}$ and $\mathcal{Q}_{abcd}$:*

$$\mathcal{S}_{abc} = 0 \iff \mathcal{C}_{abcd} = L \mathcal{Q}_{abcd}$$

where L is a complex function.

As seen above in (25), the vanishing of $\mathcal{S}'_{abcd}$ is equivalent to the proportionality between $\mathcal{C}_{abcd}$ and $\mathcal{Q}_{abcd}$ whenever there exists a potential σ for the Ernst one-form —so that the Mars-Simon tensor can be defined—. Thus, Lemma 1 implies the existence of a precise relation between the tensors $\mathcal{S}_{abc}$ and $\mathcal{S}_{abcd}$. This relation must be invariant under the freedom (13), because so is $\mathcal{S}_{abc}$, and given that $\mathcal{S}_{abcd}$ changes according to (23) this relation will not involve $\mathcal{Q}_{abcd}$. This leads to exactly the same relation (26) but simply replacing $\mathcal{C}_{abcd}$ by $\mathcal{S}_{abcd}$:

$$\mathcal{S}_{abc} = \gamma_a [b \mathcal{S}_c]_{mrd} \zeta^m \mathcal{F}^{rd} + 4 \zeta^m \zeta^r \mathcal{S}_{mar} [c \sigma_b] \quad (30)$$

The main relevance of the space-time Simon tensor resides in the following result (lemma 4 in [21])

Proposition 1. *Let $(\mathcal{M}, g_{ab})$ be a Ricci-flat, non-locally-flat, C^3 spacetime possessing a Killing vector $\vec{\zeta}$ which is non-null on a dense subset of $\mathcal{M}$ and assume that the corresponding spacetime Simon tensor vanishes everywhere. Assume also that $\mathcal{F} \cdot \mathcal{F} \neq 0$ everywhere. Then, the Ernst 1-form is exact $\sigma_a = \nabla_a \sigma$ and furthermore:*

$$\mathcal{C}_{abcd} = \frac{-1}{\sigma + a} \mathcal{Q}_{abcd}, \quad \mathcal{F} \cdot \mathcal{F} = b(\sigma + a)^4$$

where a and b are complex constants.

Combining this result with Theorem 2 we obtain another version of the local characterization of the Kerr space-time.

Theorem 3. *Under the same hypothesis of Proposition 1, $(\mathcal{M}, g_{ab})$ is locally isometric to the Kerr space-time if and only if the constant a is purely imaginary and b is real and negative.*

4 The superenergy of $\mathcal{S}_{abcd}$

Given a real Weyl candidate W_{abcd} , we define its Bel-Robinson tensor $T_{abcd}\{W\}$ by

$$T_{abcd}\{W\} \equiv W_a^p{}_d^m W_{bpcm} + W_a^m{}_c^p W_{bmdp} - \frac{1}{8} g_{ab} g_{cd} W_{mnp l} W^{mnp l}. \quad (31)$$

As described in the general framework of [32], this tensor is the basic *superenergy tensor* of the Weyl candidate W_{abcd} in 4 dimensions. We prefer to call this tensor the Bel-Robinson tensor of W_{abcd} in analogy with the Bel-Robinson tensor constructed out of the Weyl tensor [3, 32].

In the present context it is more convenient to re-write (31) in terms of the self-dual tensor $\mathcal{W}_{abcd} = W_{abcd} + iW_{abcd}^*$ corresponding to the Weyl candidate and its dual. In this way one can prove that the tensor (31) can be rewritten simply as (see e.g. [28])

$$T_{abcd}\{\mathcal{W}\} \equiv \mathcal{W}_a^p{}_c^m \overline{\mathcal{W}}_{bpd m} = T_{abcd}\{W\}. \quad (32)$$

As any other superenergy tensor, the Bel-Robinson tensor of a Weyl candidate has the following properties —see [32] for detailed proofs.

Theorem 4. If $T_{abmn}\{W\}$ is the Bel-Robinson tensor of the Weyl candidate W_{abmn} then

1. $T_{(abmn)}\{W\} = T_{abmn}\{W\}$, $T^a_{amn}\{W\} = 0$.
2. Generalized dominant property: if $\vec{u}_1, \vec{u}_2, \vec{u}_3, \vec{u}_4$ are causal future-directed vectors then $T_{abmn}\{W\}u_1^a u_2^b u_3^m u_4^n \geq 0$. The inequality is strict if $\vec{u}_1, \vec{u}_2, \vec{u}_3, \vec{u}_4$ are timelike.
3. $T_{abmn}\{W\} = 0 \iff W_{abmn} = 0 \iff W_{abmn} = 0 \iff \exists$ a set of timelike vectors $\vec{u}_1, \vec{u}_2, \vec{u}_3$ and $\vec{u}_4$ such that $T_{abmn}\{W\}u_1^a u_2^b u_3^m u_4^n = 0 \iff$ there is a timelike vector $\vec{u}$ such that $T_{abmn}\{W\}u^a u^b u^m u^n = 0$.

The last point of this theorem enables us to write the vanishing of the self-dual Weyl candidate W_{abcd} as a single scalar condition. For an arbitrary timelike vector $\vec{u}$ we define the scalar

$$U_{\vec{u}}(\mathcal{W}) \equiv T_{abmn}\{W\}u^a u^b u^m u^n \quad (33)$$

which, if $W_{abcd} \neq 0$, is a positive quantity according to points 2 and 3 of Theorem 4. Therefore one has a natural scalar quantity which enables us to measure, for each timelike vector, the proximity to the geometric conditions determined by the tensor condition $W_{abcd} = 0$. The scalar quantity $U_{\vec{u}}(\mathcal{W})$ is called the *superenergy density* of $T_{abcd}\{W\}$ with respect to $\vec{u}$ [32]² and it depends in general on the timelike vector $\vec{u}$.

However, in those cases where there is a timelike vector defined invariantly one can select a superenergy density that acquires an invariant meaning. We can apply this idea to the particular case in which $W_{abcd} = \mathcal{S}_{abcd}$. At those points of $\mathcal{M}$ in which the Killing vector $\vec{\zeta}$ is timelike the scalar $U_{\vec{\zeta}}(\mathcal{S})$ is positive and it will vanish if and only if $\mathcal{S}_{abcd} = 0$. Given that $\vec{\zeta}$ is defined invariantly we can take the quantity $U_{\vec{\zeta}}(\mathcal{S})$ as a local invariant measure of the deviation of the spacetime $(\mathcal{M}, g_{ab})$ to the geometric conditions entailed by $\mathcal{S}_{abcd} = 0$.

Theorem 5.

$$U_{\vec{\zeta}}(\mathcal{S}) = \mathcal{E}_{ab}\bar{\mathcal{E}}^{ab} + \frac{3}{4|\sigma|^2} (3(\sigma \cdot \bar{\sigma})^2 - |\sigma \cdot \sigma|^2 + 4\text{Re}(\sigma \mathcal{E}_{ab}\bar{\sigma}^a \bar{\sigma}^b)), \quad (34)$$

where

$$\mathcal{E}_{ab} \equiv \mathcal{C}_{apbq}\zeta^p \zeta^q, \quad \sigma \cdot \sigma \equiv \sigma_a \sigma^a, \quad \sigma \cdot \bar{\sigma} \equiv \sigma_a \bar{\sigma}^a. \quad (35)$$

Proof. Starting from (15) and using the definition of the Ernst 1-form, a straightforward computation yields

$$\mathcal{U}_{ac} \equiv \mathcal{S}_{abcd}\zeta^b \zeta^d = \mathcal{E}_{ac} + \frac{\mathcal{F} \cdot \mathcal{F}}{2\sigma} \gamma_{ac} + \frac{3\sigma_a \sigma_c}{2\sigma}. \quad (36)$$

Next, we write the definition of $U_{\vec{\zeta}}(\mathcal{S})$ adapting the definition of $T_{abcd}\{\mathcal{S}\}$ to eq. (32). The result is

$$U_{\vec{\zeta}}(\mathcal{S}) = \mathcal{U}_{ac}\bar{\mathcal{U}}^{ac}. \quad (37)$$

Now we need to replace in this expression the value of $\mathcal{U}_{ab}$ found above and work out the resulting expression. After a computation, we find

$$\mathcal{U}_{ac}\bar{\mathcal{U}}^{ac} = \mathcal{E}_{ab}\bar{\mathcal{E}}^{ab} + \frac{3}{2\sigma}\bar{\mathcal{E}}_{ab}\sigma^a \sigma^b + \frac{3}{2\bar{\sigma}}\mathcal{E}_{ab}\bar{\sigma}^a \bar{\sigma}^b + \frac{3}{4|\sigma|^2} \left(3(\sigma \cdot \bar{\sigma})^2 - \lambda(\bar{\mathcal{F}} \cdot \bar{\mathcal{F}} \sigma \cdot \sigma + \mathcal{F} \cdot \mathcal{F} \bar{\sigma} \cdot \bar{\sigma}) + \lambda^2 |\mathcal{F} \cdot \mathcal{F}|^2 \right). \quad (38)$$

The final expression (34) follows after using in this expression the identity

$$\lambda \mathcal{F} \cdot \mathcal{F} = \sigma \cdot \sigma, \quad (39)$$

which is a consequence of (14). $\square$

Remark 2. Note that the result is valid for a Killing vector $\vec{\zeta}$ of any causal character.

²The notation differs from that of [32].

One may also use the previous definitions and results to introduce the quantity

$$U_{\vec{\zeta}}(\mathcal{Q}) \equiv T_{abcd}\{\mathcal{Q}\}\zeta^a\zeta^b\zeta^c\zeta^d \quad (40)$$

for the Weyl candidate (22). According to Theorem 4, the scalar $U_{\vec{\zeta}}(\mathcal{Q})$ will be non-negative whenever $\vec{\zeta}$ is causal. Indeed a straightforward computation yields

$$U_{\vec{\zeta}}(\mathcal{Q}) = \frac{9(\boldsymbol{\sigma} \cdot \bar{\boldsymbol{\sigma}})^2 - 3|\boldsymbol{\sigma} \cdot \boldsymbol{\sigma}|^2}{4}. \quad (41)$$

Hence (34) can be rendered in the form

$$U_{\vec{\zeta}}(\mathcal{S}) = U_{\vec{\zeta}}(\mathcal{C}) + \frac{U_{\vec{\zeta}}(\mathcal{Q})}{|\boldsymbol{\sigma}|^2} + \frac{3\text{Re}(\sigma\mathcal{E}_{ab}\bar{\sigma}^a\bar{\sigma}^b)}{|\boldsymbol{\sigma}|^2}. \quad (42)$$

4.1 Several alternative expressions for the scalar $U_{\vec{\zeta}}(\mathcal{S})$

The superenergy construction can be carried out for arbitrary tensors of any rank, and not only for the Weyl candidates [32]. For a 2-form F_{ab} the corresponding superenergy tensor $T_{ab}\{F\}$ is simply the standard energy-momentum tensor of a Maxwell field and can be expressed in any one of the following equivalent forms

$$T_{ab}\{F\} \equiv t_{ab} = F_{ac}F_b{}^c - \frac{1}{4}g_{ab}F_{cd}F^{cd} = \frac{1}{2}(F_{ac}F_b{}^c + F_{ac}^*F_b^{*c}) = \frac{1}{2}\mathcal{F}_{ac}\bar{\mathcal{F}}_b{}^c. \quad (43)$$

Using these equations and (9) we can easily derive the identities

$$t_{ab} = t_{ba}, \quad t^a{}_a = 0, \quad t_{ac}t_b{}^c = \frac{1}{4}g_{ab}t_{cd}t^{cd}, \quad t_{cd}t^{cd} = \frac{1}{16}(\mathcal{F} \cdot \mathcal{F})(\bar{\mathcal{F}} \cdot \bar{\mathcal{F}}),$$

and by virtue of the ‘‘Maxwell’’ equations $\nabla^a\mathcal{F}_{ab} = 0$, t_{ab} is divergence free

$$\nabla^a t_{ab} = 0. \quad (44)$$

As any other superenergy tensor, it also possesses the dominant property, that is to say, $t_{ab}u_1^a u_2^b \geq 0$ for arbitrary future-pointing vectors $\vec{u}_1$ and $\vec{u}_2$, and for timelike $\vec{u}$ one has

$$t_{ab}u^a u^b \geq 0, \quad t_{ab}u^a u^b = 0 \iff t_{ab} = 0 \iff F_{ab} = 0.$$

On using Eq.(14) the tensor t_{ab} can be written in our case as

$$t_{ab} = \frac{1}{8\lambda^2} [2(\boldsymbol{\sigma} \cdot \bar{\boldsymbol{\sigma}})\zeta_a\zeta_b + \lambda(\sigma_a\bar{\sigma}_b + \bar{\sigma}_a\sigma_b) - \lambda(\boldsymbol{\sigma} \cdot \bar{\boldsymbol{\sigma}})g_{ab} + 2\zeta_{(a}v_{b)}] \quad (45)$$

where we have introduced the real one-form

$$v_b \equiv i\eta_{bcde}\bar{\sigma}^c\zeta^d\sigma^e$$

which obviously satisfies

$$v_b\zeta^b = v_b\sigma^b = v_b\bar{\sigma}^b = 0, \quad v_b v^b = \lambda [(\boldsymbol{\sigma} \cdot \boldsymbol{\sigma})(\bar{\boldsymbol{\sigma}} \cdot \bar{\boldsymbol{\sigma}}) - (\boldsymbol{\sigma} \cdot \bar{\boldsymbol{\sigma}})^2].$$

With these definitions at hand, we can find an interesting expression for $\mathcal{E}_{ac}$. To that end, simply use the first in (10) to compute

$$\mathcal{E}_{ac} = -\zeta^b\nabla_c\mathcal{F}_{ab} = -\frac{1}{2}\nabla_c\sigma_a + \mathcal{F}_{ab}\nabla_c\zeta^b = -\frac{1}{2}\nabla_c\sigma_a + \frac{1}{2}\mathcal{F}_{ab}(\mathcal{F}_c{}^b + \bar{\mathcal{F}}_c{}^b)$$

so that introducing here the identity (9) together with (39) and the definition of t_{ab} we arrive at the desired expression

$$\mathcal{E}_{ac} = t_{ac} + \frac{1}{8\lambda}(\boldsymbol{\sigma} \cdot \boldsymbol{\sigma})g_{ac} - \frac{1}{2}\nabla_c\sigma_a. \quad (46)$$

Observe that this provides a very simple formula for the electric and magnetic parts of the Weyl tensor with respect to the preferred timelike direction $\vec{\zeta}$ in Ricci-flat stationary spacetimes in terms of the Ernst one-form.

Using (46), (27) and (36) we can also derive

$$\mathcal{U}_{ac} = t_{ac} + \frac{1}{8\lambda}(\boldsymbol{\sigma} \cdot \boldsymbol{\sigma})g_{ac} - \frac{1}{2}\nabla_c \sigma_a + \frac{(\boldsymbol{\sigma} \cdot \boldsymbol{\sigma})}{2\lambda\sigma}\gamma_{ac} + \frac{3\sigma_a\sigma_c}{2\sigma}. \quad (47)$$

From the previous formula (46) it is very easy to compute the Bel-Robinson superenergy with respect to the direction $\vec{\zeta}$:

$$U_{\vec{\zeta}}(\mathcal{C}) = \frac{1}{4} [\nabla_a \sigma_b \nabla^a \bar{\sigma}^b - 2t_{ab} \nabla^a (\sigma^b + \bar{\sigma}^b)] = \frac{1}{4} [\nabla_a \nabla_b \sigma \nabla^a \nabla^b \bar{\sigma} - 4t_{ab} \nabla^a \nabla^b \lambda]$$

which on using (45) can also be written in terms of the Ernst one-form exclusively as

$$U_{\vec{\zeta}}(\mathcal{C}) = \frac{1}{4} \nabla_a \sigma_b \nabla^a \bar{\sigma}^b - \frac{1}{16\lambda^2}(\boldsymbol{\sigma} \cdot \boldsymbol{\sigma})(\bar{\boldsymbol{\sigma}} \cdot \bar{\boldsymbol{\sigma}}) + \frac{(\boldsymbol{\sigma} \cdot \bar{\boldsymbol{\sigma}})}{32\lambda^2} [(\boldsymbol{\sigma} \cdot \boldsymbol{\sigma}) + (\bar{\boldsymbol{\sigma}} \cdot \bar{\boldsymbol{\sigma}})] - \frac{1}{8\lambda} \bar{\sigma}^a \sigma^b \nabla_a (\sigma_b + \bar{\sigma}_b). \quad (48)$$

This is an expression for the Bel-Robinson super-energy relative to the direction $\vec{\zeta}$ involving only the Ernst one-form and its derivatives.

We can also use (46) together with (43) to compute

$$\mathcal{E}_{ab} \bar{\sigma}^a \bar{\sigma}^b = \frac{(\bar{\boldsymbol{\sigma}} \cdot \bar{\boldsymbol{\sigma}})}{8\lambda} [(\boldsymbol{\sigma} \cdot \boldsymbol{\sigma}) + (\boldsymbol{\sigma} \cdot \bar{\boldsymbol{\sigma}})] - \frac{1}{2} \bar{\sigma}^a \bar{\sigma}^b \nabla_b \sigma_a.$$

The previous two expressions together with (42-41) lead after some manipulations to

$$\begin{aligned} U_{\vec{\zeta}}(\mathcal{S}) &= \frac{1}{4} \nabla_b \bar{\sigma}_a \nabla^b \sigma^a + \frac{9(\boldsymbol{\sigma} \cdot \bar{\boldsymbol{\sigma}})^2}{4|\sigma|^2} - (6\lambda^2 - 6\lambda + |\sigma|^2) \frac{|\boldsymbol{\sigma} \cdot \boldsymbol{\sigma}|^2}{16\lambda^2|\sigma|^2} + \\ &2\text{Re} \left[\frac{(\boldsymbol{\sigma} \cdot \bar{\boldsymbol{\sigma}})(\boldsymbol{\sigma} \cdot \boldsymbol{\sigma})}{32\lambda^2|\sigma|^2} (6\lambda + \sigma) \bar{\sigma} - \frac{\sigma^a \bar{\sigma}^b}{8\lambda} \nabla_b \sigma_a - \frac{3\sigma^a \sigma^b}{4\sigma} \nabla_b \bar{\sigma}_a \right], \end{aligned} \quad (49)$$

which again is an expression depending exclusively on the Ernst potential and its derivatives.

We can summarize the previous results in the following theorem

Theorem 6. *Let $(\mathcal{M}, g_{ab})$ be a smooth spacetime with a Killing vector $\vec{\zeta}$ that contains a stationary asymptotically flat 4-end $\mathcal{M}_\infty$ such that $\vec{\zeta}$ tends to a time translation at infinity and the Komar mass of $\vec{\zeta}$ at $\mathcal{M}_\infty$ does not vanish. Fix the freedom (13) such that $\sigma \rightarrow 0$ at $\mathcal{M}_\infty$.*

Then, the scalar (49) constructed exclusively from σ and its first and second derivatives is non-negative on the entire region where $\vec{\zeta}$ is timelike, and it vanishes if and only if $(\mathcal{M}, g_{ab})$ is locally isometric to the Kerr spacetime.

5 Quality factors that measure the Kerr-ness of stationary spacetimes

In this section we present several quality factors to measure how close a given spacetime is to the exterior part of the Kerr solution. By a quality factor we mean an adimensional scalar function q with $q \in [0, 1]$ and such that its value provides a measure of the “Kerr-ness” of the given space-time at any point.

We present three different possibilities. All of them can be applied to generic stationary spacetimes, the first two to stationary Ricci-flat ones, while the first one can only be used when the space-time is Ricci-flat and has an asymptotically flat end.

5.1 A quality factor for stationary Ricci-flat asymptotically flat spacetimes

Theorem 6 provides a neat characterization for the Kerr solution in terms of a single scalar and real quantity, which happens to be strictly positive if the metric is not Kerr. One can thus think that the scalar (49) may provide a measure of how much a region of the spacetime differs from the stationary portion of Kerr spacetime: the smaller the scalar (49) is, the better resemblance to the Kerr solution. Unfortunately, the previous sentence does not have an absolute meaning, because the scalar (49) has physical units —of $(\text{length})^{-4}$ — and thus we can always choose the physical dimensions such that it becomes large or small.

What is needed here is another similar quantity that normalizes (49) and eliminates the physical units. There may be several choices here, but our own, which we believe is a natural one, is based again on the superenergy construction and its positivity properties. As a matter of fact, one can define a (positive-definite and sesquilinear) inner product $\langle \cdot, \cdot \rangle_{\bar{\zeta}}$ on the complex vector space of self-dual Weyl candidates as follows:

$$\langle \mathcal{W}_1, \mathcal{W}_2 \rangle_{\bar{\zeta}} \equiv \mathcal{W}_{1a}{}^p{}_c{}^m \bar{\mathcal{W}}_{2bpdm} \zeta^a \zeta^b \zeta^c \zeta^d$$

Sesquilinearity follows obviously from the definition, while its positive definiteness is a direct consequence of Theorem 4. Observe that the norm here is simply the superenergy scalar introduced in (33)

$$\| \mathcal{W} \|_{\bar{\zeta}}^2 \equiv \langle \mathcal{W}, \mathcal{W} \rangle_{\bar{\zeta}} = U_{\bar{\zeta}}(\mathcal{W}). \quad (50)$$

Definition 1 (Quality factor q). *The quality factor measuring the Kerr-ness of any stationary region in a non-flat asymptotically flat and Ricci-flat spacetime $(\mathcal{M}, g_{ab})$ is defined by*

$$q = 1 - Q^2$$

where

$$Q^2 \equiv \frac{U_{\bar{\zeta}}(\mathcal{S})}{\left(\sqrt{U_{\bar{\zeta}}(\mathcal{C})} + |\sigma|^{-1} \sqrt{U_{\bar{\zeta}}(\mathcal{Q})} \right)^2}.$$

The main properties of this quality factor are collected in the following

Theorem 7. *The quality factor q has the following properties:*

1. $q \in [0, 1]$.
2. $q = 1$ on an open neighborhood $U \subset \mathcal{M}$ if and only if (U, g_{ab}) is locally isometric to the Kerr solution.
3. At any point $x \in \mathcal{M}$, $q = 0$ if and only if $\mathcal{C}_{abcd}|_x = k(\mathcal{Q}_{abcd}/\sigma)|_x$ for a positive real constant $k \in \mathbb{R}^+$.

Remark 3. Notice that the condition for $q = 0$ is that the space-time is of Petrov type D —as it is the case for the Kerr solution—, however the proportionality constant k has the *opposite* sign to the case of Kerr.

Proof. From (50) we can rewrite Q^2 as

$$Q^2 = \frac{\| \mathcal{S} \|_{\bar{\zeta}}^2}{\left(\| \mathcal{C} \|_{\bar{\zeta}} + \left\| \frac{\mathcal{Q}}{\sigma} \right\|_{\bar{\zeta}} \right)^2}$$

and using here that (indices suppressed) $\mathcal{S} = \mathcal{C} + \mathcal{Q}/\sigma$ the triangle inequality $\| \mathcal{C} + \mathcal{Q}/\sigma \|_{\bar{\zeta}} \leq \| \mathcal{C} \|_{\bar{\zeta}} + \| \mathcal{Q}/\sigma \|_{\bar{\zeta}}$ leads to

$$0 \leq Q^2 \leq 1$$

which proves 1. Now, Theorem 6 informs us that the space-time is locally isometric to Kerr if and only if $U_{\vec{\zeta}}(\mathcal{S}) = \|\mathcal{S}\|_{\vec{\zeta}}^2 = 0$, that is, if $Q^2 = 0$, which proves 2. To prove 3, recall that the triangle inequality is an equality if and only if $\mathcal{Q}/\sigma|_x$ and $\mathcal{C}|_x$ are *positively* linearly dependent, i.e. if $\mathcal{C}|_x = k(\mathcal{Q}/\sigma)|_x$ with $k \geq 0$. The case with $k = 0$ is excluded as the space-time would be flat. $\square$

Remark 4. It is very important to remark that q can be computed in any given non-flat Ricci-flat and asymptotically flat stationary space-time *without any reference* to the Kerr solution. One does not need to map the space-time into the Kerr solution, nor is it necessary to compare or choose coordinates relative to Kerr. All that one needs is an explicit expression of the metric of the space-time to be compared to the Kerr solution. Notice, furthermore, that using (49), (48) and (41) one can immediately write an expression for q in terms of σ and its derivatives.

At any point x of the stationary region in $(\mathcal{M}, g_{ab})$ one can say that x is “ $(q \times 100)\%$ Kerr” (so it is 100% Kerr if $q = 1$, and 0% Kerr if $q = 0$). The precise value of q which gives a good approximation to Kerr depends on the context, and it will have to be calibrated by studying several examples. Some of these are given in Section 6.

5.2 Quality factors for stationary Ricci-flat spacetimes

There may arise situations where one does not know if the entire space-time is asymptotically flat, or also when one knows that there are no asymptotically flat ends but still the proximity with a Kerr space-time is suspected, for example close to the ergo-surface of the Killing vector or to a possible null hypersurface which one suspects may resemble the horizon of Kerr space-time. In these situations we cannot apply the previous quality factor, which requires fixing the constant α in (13) for the potential σ . However, we can still make use of the previous results by complicating things a little bit, as follows.

For Ricci-flat spacetimes we can make use of Theorems 2 and 3 which do not require asymptotic flatness. The first thing we must require to use those theorems, according to (25) or Lemma 1, is the proportionality between $\mathcal{C}_{abcd}$ and $\mathcal{Q}_{abcd}$. Note, however, that this can be easily measured by means of

$$q_1 \equiv \frac{|\langle \mathcal{C}, \mathcal{Q} \rangle_{\vec{\zeta}}|^2}{\|\mathcal{C}\|_{\vec{\zeta}}^2 \|\mathcal{Q}\|_{\vec{\zeta}}^2}.$$

Theorem 8. *The quality factor q_1 has the following properties*

1. $q_1 \in [0, 1]$.
2. $q_1 = 1$ at a point $x \in \mathcal{M}$ if and only if $\exists l \in \mathbb{C} : \mathcal{C}_{abcd}|_x = l \mathcal{Q}_{abcd}|_x$.
3. $q_1 = 0$ at a point $x \in \mathcal{M}$ if and only if $\mathcal{E}_{ab} \bar{\sigma}^a \bar{\sigma}^b|_x = 0$.

Proof. The Cauchy-Schwarz inequality immediately gives

$$0 \leq q_1 \leq 1, \quad q_1|_x = 1 \iff \mathcal{C}_{abcd}|_x = l \mathcal{Q}_{abcd}|_x, \quad (51)$$

for some $l \in \mathbb{C}$. The other extreme value, $q_1 = 0$, can only occur when

$$\langle \mathcal{C}, \mathcal{Q} \rangle_{\vec{\zeta}} = \frac{3}{2} \mathcal{E}_{ab} \bar{\sigma}^a \bar{\sigma}^b$$

vanishes, proving point 3. $\square$

Point 2 in this theorem informs us that q_1 measures how much $\mathcal{C}_{abcd}$ and $\mathcal{Q}_{abcd}$ are proportional to each other. A possible alternative to measure this proportionality is the use of the space-time Simon tensor $\mathcal{S}_{abc}$. We first note the following simple result.

Proposition 2. *Assume that the Killing vector $\vec{\zeta}$ is time-like. Then*

$$\mathcal{S}_{abc} \bar{\mathcal{S}}^{abc} \geq 0, \quad \mathcal{S}_{abc} \bar{\mathcal{S}}^{abc} = 0 \iff \mathcal{S}_{abc} = 0. \quad (52)$$

Proof. Since $\vec{\zeta}$ is time-like, from (29) we know that $\mathcal{S}_{abc}$ is totally orthogonal to $\vec{\zeta}$, and thus spatial with respect to the Killing vector, from where the result follows easily. $\square$

This result together with Lemma 1 enables us to use the quantity $\mathcal{S}_{abc}\overline{\mathcal{S}}^{abc}$ in a definition of the quality factor to measure the proportionality of $\mathcal{C}_{abcd}$ and $\mathcal{Q}_{abcd}$. To that end, for any two Lanczos candidates $\mathcal{S}_{abc}^1$ and $\mathcal{S}_{abc}^2$ (not necessarily traceless), define

$$\langle \mathcal{S}^1, \mathcal{S}^2 \rangle \equiv \mathcal{S}_{abc}^1 \overline{\mathcal{S}}^{2abc}. \quad (53)$$

This is again a sesquilinear product on the space of Lanczos candidates (not necessarily traceless), and it is positive definite when acting on candidates that are totally orthogonal to a timelike Killing vector $\vec{\zeta}$, as follows from Proposition 2. Actually,

$$\| \mathcal{S} \|^2 \equiv \langle \mathcal{S}, \mathcal{S} \rangle$$

happens to be essentially the superenergy of the (real and imaginary parts) of $\mathcal{S}_{abc}$ with respect to $\vec{\zeta}$, according to the general construction in [32].

Now, observe that $\mathcal{S}_{abc}$ can be written, from its definition (26), directly as the sum $\mathcal{S}_{abc}^1 + \mathcal{S}_{abc}^2$ where

$$\mathcal{S}_{abc}^1 \equiv -\frac{1}{2}\gamma_{a[b}\nabla_{c]}(\mathcal{F} \cdot \mathcal{F}), \quad \mathcal{S}_{abc}^2 \equiv -4\mathcal{E}_{a[b}\sigma_{c]},$$

which are Lanczos candidates on their own, and they are totally orthogonal to $\vec{\zeta}$. Therefore we can define

$$q_2 \equiv 1 - \frac{\| \mathcal{S} \|^2}{(\| \mathcal{S}^1 \| + \| \mathcal{S}^2 \|)^2}. \quad (54)$$

Theorem 9. *The quality factor q_2 has the following properties*

1. $q_2 \in [0, 1]$.
2. $q_2 = 1$ at a point $x \in \mathcal{M}$ if and only if $\exists l \in \mathbb{C} : \mathcal{C}_{abcd}|_x = l\mathcal{Q}_{abcd}|_x$.
3. $q_2 = 0$ at a point $x \in \mathcal{M}$ if and only if $\nabla_c(\mathcal{F} \cdot \mathcal{F})|_x = 0$ (thus $\mathcal{F} \cdot \mathcal{F} = \text{const.}$ if $q_2 = 0$ on an open neighborhood) which entails $\mathcal{S}_{abc}^1|_x = 0$.

Proof. The first property is a direct consequence of the triangle inequality $\| \mathcal{S} \| = \| \mathcal{S}^1 + \mathcal{S}^2 \| \leq \| \mathcal{S}^1 \| + \| \mathcal{S}^2 \|$, while $q_2|_x = 1$ happens if and only if $\mathcal{S}_{abc}|_x = 0$ as is clear from (54), so that Lemma 1 entails then $\mathcal{C}_{abcd}|_x = l\mathcal{Q}_{abcd}|_x$ for some $l \in \mathbb{C}$. To prove the third point, recall that $q_2|_x = 0$ if and only if $\mathcal{S}_{abc}^1|_x$ and $\mathcal{S}_{abc}^2|_x$ are positively proportional to each other, that is

$$\mathcal{S}_{abc}^1|_x = k\mathcal{S}_{abc}^2|_x, \quad k \in \mathbb{R}, \quad k \geq 0$$

which reads

$$\frac{1}{2}\gamma_{a[b}\nabla_{c]}(\mathcal{F} \cdot \mathcal{F})|_x = 4k\mathcal{E}_{a[b}\sigma_{c]}|_x.$$

Contracting here b and a we derive

$$\lambda\nabla_c(\mathcal{F} \cdot \mathcal{F})|_x = 4k\mathcal{E}_{ac}\sigma^a|_x$$

but using here (46) and (45) together with (39) we also have

$$\mathcal{E}_{ac}\sigma^a = \frac{\sigma \cdot \sigma}{8\lambda}(\sigma_c + \bar{\sigma}_c) - \frac{1}{2}\sigma^b\nabla_c\sigma_b = \frac{\sigma \cdot \sigma}{4\lambda}\nabla_c\lambda - \frac{1}{4}\nabla_c(\sigma \cdot \sigma) = -\frac{\lambda}{4}\nabla_c(\mathcal{F} \cdot \mathcal{F}) \quad (55)$$

so that the previous condition becomes

$$(1+k)\lambda\nabla_c(\mathcal{F} \cdot \mathcal{F})|_x = 0$$

which proves the result. $\square$

Actually, eq.(55) allows us to derive an alternative useful expression for $\mathcal{S}_{abc}$.

Lemma 2. *The space-time Simon tensor can be written in the form*

$$\mathcal{S}_{abc} = \frac{1}{\lambda} (4\mathcal{E}_{a[b}\gamma_{c]d} + 2\mathcal{E}_{d[c}\gamma_{b]a})\sigma^d. \quad (56)$$

Proof. By noting that $\gamma_{cd}\sigma^d = -\lambda\sigma_c$ the first summand provides directly $\mathcal{S}_{abc}^2$ while the second summand is $\mathcal{S}_{abc}^1$ as follows from (55). $\square$

If we combine eq. (51) with the second point of Theorem 9 we deduce that

$$q_1 = 1 \iff q_2 = 1. \quad (57)$$

One can find an alternative, useful, expression for q_2 as follows. By explicit computation one can easily obtain the following results

$$\|\mathcal{S}^1\|^2 = \frac{\lambda^2}{4} \nabla_c (\mathcal{F} \cdot \mathcal{F}) \nabla^c (\overline{\mathcal{F}} \cdot \overline{\mathcal{F}}) \quad (58)$$

$$\langle \mathcal{S}^1, \mathcal{S}^2 \rangle = -\|\mathcal{S}^1\|^2 \quad (59)$$

$$\|\mathcal{S}^2\|^2 = 8 [(\sigma \cdot \bar{\sigma}) \mathcal{E}_{ab} \bar{\mathcal{E}}^{ab} - \mathcal{E}_b{}^c \bar{\mathcal{E}}_{ac} \sigma^a \bar{\sigma}^b] \quad (60)$$

where we have used (55) and (58) to get (59). From these one immediately finds

$$\langle \mathcal{S}^1, \mathcal{S} \rangle = 0, \quad \|\mathcal{S}\|^2 = \|\mathcal{S}^2\|^2 - \|\mathcal{S}^1\|^2 \quad (61)$$

which also implies

$$\|\mathcal{S}^2\|^2 \geq \|\mathcal{S}^1\|^2.$$

Observe that the second in (61) allows us to rewrite q_2 as given in (54) in a simpler form

$$q_2 = 1 - \frac{\|\mathcal{S}^2\|^2 - \|\mathcal{S}^1\|^2}{(\|\mathcal{S}^1\| + \|\mathcal{S}^2\|)^2} = 1 - \frac{\|\mathcal{S}^2\| - \|\mathcal{S}^1\|}{\|\mathcal{S}^1\| + \|\mathcal{S}^2\|} = \frac{2\|\mathcal{S}^1\|}{\|\mathcal{S}^1\| + \|\mathcal{S}^2\|}$$

from where the properties in Theorem 9 become more transparent.

Remark 5. We note that the quality factors q_1 and q_2 do not involve the Ernst potential and hence they can be computed for any space-time possessing a Killing vector, be it Ricci flat or not.

Assume then that any of q_1 or q_2 were unity. In that case, from Proposition 1 we know that, for Ricci-flat spacetimes, there would exist a complex constant a such that L above would be $L = -1/(\sigma + a)$ for a potential σ with $\text{Re}(\sigma) = 1 + \lambda$ (redefining a if necessary). Then we would have

$$\langle \mathcal{C}, \mathcal{Q} \rangle_{\bar{\xi}} = -\frac{1}{\sigma + a} \|\mathcal{Q}\|_{\bar{\xi}}^2 \quad (62)$$

so that the Kerr condition $a = -\bar{a}$ (see Theorem 3) would require that the quantity

$$\beta \equiv \text{Re} \left(\frac{\|\mathcal{Q}\|_{\bar{\xi}}^2}{\langle \mathcal{C}, \mathcal{Q} \rangle_{\bar{\xi}}} \right)$$

equals $-(1 + \lambda)$. Notice once again that to compute β we only need to know the space-time and the Killing vector, and therefore we can control this property by means of the partial quality factor

$$\kappa \equiv 2 \frac{|\beta + 1||\lambda| - (\beta + 1)\lambda}{(|\beta + 1| + |\lambda|)^2} = -2\lambda \frac{|\beta + 1| + (\beta + 1)}{(|\beta + 1| - \lambda)^2} \quad (63)$$

where in the second equality we have used that $\lambda < 0$. It is easy to prove that

Proposition 3. *The quantity κ fulfills the following properties*

- $0 \leq \kappa \leq 1$,
- $\kappa = 0$ if and only if $\beta + 1 < 0$,

- $\kappa = 1$ if and only if $\beta + 1 = -\lambda$ (this is the Kerr condition).

Proof. An alternative expression for κ is

$$\kappa = 1 - \frac{(\beta + 1 + \lambda)^2}{(|\beta + 1| + |\lambda|)^2}$$

from where the first and third properties are obvious. The second follows directly from (63). $\square$

Provided that $q_1 = 1$ (or equivalently $q_2 = 1$) we must finally control the second condition on Theorem 3, that is to say that the constant b in Proposition 1 be real and negative. From (62) and (39) this can be rewritten as

$$z \equiv (\boldsymbol{\sigma} \cdot \boldsymbol{\sigma}) \frac{\langle \mathcal{C}, \mathcal{Q} \rangle_{\bar{z}}^4}{\|\mathcal{Q}\|_{\bar{z}}^8}$$

being real and positive (as this would have to be $z = b\lambda$). Then set

$$\varkappa \equiv 1 - \frac{1}{4} \left(\frac{z}{|z|} - 1 \right) \left(\frac{\bar{z}}{|z|} - 1 \right).$$

Proposition 4. *The following statements hold true*

- $0 \leq \varkappa \leq 1$,
- $\varkappa = 0$ if and only if $z = -|z| = \bar{z}$ (i.e. z is real and negative).
- $\varkappa = 1/2$ if and only if $z = -\bar{z}$ (i.e. z is purely imaginary).
- $\varkappa = 1$ if and only if $z = |z| = \bar{z}$ (i.e. z is real and positive). This is the Kerr condition.

Proof. If we set $z/|z| = e^{i\phi}$ then elementary manipulations show that the scalar $\varkappa$ takes the following form in terms of ϕ

$$\varkappa = \cos^2 \left(\frac{\phi}{2} \right), \quad (64)$$

which makes it clear that $0 \leq \varkappa \leq 1$. Now, one has

- $\varkappa = 0 \iff \phi = (2n - 1)\pi, n \in \mathbb{Z} \iff z$ is real and negative.
- $\varkappa = 1/2 \iff \phi = \pi/2 + n\pi, n \in \mathbb{Z} \iff z$ is purely imaginary.
- $\varkappa = 1 \iff \phi = 2n\pi, n \in \mathbb{Z} \iff z$ is real and positive.

$\square$

Thus, we can define two new quality factors to measure the Kerr-ness of general stationary Ricci-flat spacetimes.

Definition 2 (Quality factors $\hat{q}$ and $\tilde{q}$). *Two quality factors measuring the Kerr-ness of any stationary region in a Ricci-flat spacetime $(\mathcal{M}, g_{ab})$ are defined by*

$$\hat{q} \equiv q_2 \kappa \varkappa, \quad \tilde{q} \equiv q_1 \kappa \varkappa,$$

The main properties of these quality factors are

Theorem 10. *The quality factors $\hat{q}$ and $\tilde{q}$ have the following properties:*

1. $\hat{q}, \tilde{q} \in [0, 1]$.
2. On an open neighborhood $U \subset \mathcal{M}$ we have that $\hat{q} = 1 \iff \tilde{q} = 1 \iff (U, g_{ab})$ is locally isometric to the Kerr solution.
3. At any point $x \in \mathcal{M}$, $\hat{q}|_x = 0$ if and only if either $(\beta + 1)|_x < 0$, or $z|_x$ is real and negative, or $\nabla_c (\mathcal{F} \cdot \mathcal{F})|_x = 0$.

4. At any point $x \in \mathcal{M}$, $\tilde{q} = 0$ if and only if either $(\beta + 1)|_x < 0$, or $z|_x$ is real and negative, or $\mathcal{E}_{ab}\bar{\sigma}^a\bar{\sigma}^b|_x = 0$.

Proof. The first item follows from the fact that each quality factor is the product of quantities in the interval $[0, 1]$. To prove the second item, we note that if $\hat{q} = 1$, then clearly $q_2 = 1$, $\kappa = 1$ and $\varkappa = 1$, as $q_2, \kappa, \varkappa \in [0, 1]$. This together with (57) implies that

$$\hat{q} = 1 \iff \tilde{q} = 1.$$

According to Proposition 4, Proposition 3, Theorem 8 and Theorem 9, the fact that $q_2 = 1$, $q_1 = 1$, $\kappa = 1$, $\varkappa = 1$ in an open neighborhood $U \subset \mathcal{M}$ is equivalent to the conditions of Theorem 3 being met and hence (U, g_{ab}) is locally isometric to the Kerr solution. Similarly, points 3 and 4 of the theorem follow directly from the mentioned propositions and theorems. $\square$

Remark 6. One could also have defined $\hat{q}' \equiv (q_2 + \kappa + \varkappa)/3$ —and the same for $\tilde{q}$ —which also has properties 1 and 2 of the theorem. However, observe that the chosen $\hat{q}$ is more demanding, in the sense that larger values of $\hat{q}$ are required to have a good resemblance to the Kerr space-time. For example, if $q_2 = 1$, $\kappa = 1/2$ and $\varkappa = 1/2$ we would have $\hat{q}' = 2/3$ while $\hat{q} = 1/4$.

5.3 How to apply the quality factors to general stationary spacetimes

The results described so far assume that $(\mathcal{M}, g_{ab})$ is a vacuum solution of the Einstein field equations (without cosmological constant Λ) and hence they cannot be used to establish the proximity of a non-Ricci-flat solution to the Kerr spacetime. The problems are of two types. First of all, the Mars-Simon tensor can only be defined on spacetimes with a potential σ for the Ernst one-form, and this only happens when $\zeta_{[b}R_{a]c}\zeta^c = 0$. A way to circumvent this is to replace the Mars-Simon tensor with another quantity which can be defined in any spacetime, independently of the existence of the Ernst potential. Obviously, the space-time Simon tensor is an useful alternative, as it can be defined in any spacetime with a Killing vector, be it Ricci-flat or not, and independently of the existence of a potential σ . And this is where the second difficulty arises, because the vanishing of the space-time Simon tensor implies the proportionality of $\mathcal{C}_{abcd}$ and $\mathcal{Q}_{abcd}$, but the explicit form of the proportionality factor given in Proposition 1, as well as the explicit expression appearing there for $\mathcal{F} \cdot \mathcal{F}$, depend crucially on the assumption that the Ricci tensor vanishes.

Nevertheless, it is not only desirable, but actually important, to be able to measure the proximity to Kerr for stationary spacetimes with non-vanishing Ricci tensor. And this is important at least in three respects: (i) to consider approximate solutions to the Einstein vacuum field equations and check their resemblance to the Kerr space-time, (ii) for cases with a cosmological constant Λ , and (iii) for more general cases such as solutions for a perfect-fluid energy-momentum tensor, or matched solutions containing a vacuum region and an interior part, and so on. In all cases we are going to argue that we can actually use some of the previous results conveniently adapted to the situation under consideration, and with some precautions.

We split this analysis in two, the case with an Ernst potential, and the case without it.

5.3.1 The case with an Ernst potential: $R_{ab}\zeta^b = \mu\zeta_a$

This case includes, obviously, the vacuum case with a cosmological constant $R_{ab} = \Lambda g_{ab}$, as well as for instance any space-time with a *rigidly rotating* fluid, which are characterized by having an energy-momentum tensor (ergo a Ricci tensor) with a timelike eigendirection proportional to a timelike Killing vector ζ .

Now the relations of Proposition 1 no longer hold in general, as they are derived using the Ricci-flat condition. Nevertheless, the factors κ and $\varkappa$ can still be defined while q_1 and q_2 still measure the proportionality between $\mathcal{C}_{abcd}$ and $\mathcal{Q}_{abcd}$. One can expect intuitively that, provided the space-time is actually close to the Kerr solution somewhere, the deviation from Ricci-flatness will be small there, and the relations of Proposition 1 will hold *approximately*, that is, with some small corrections proportional to the non-zero Ricci components. Thus, it seems clear that the quality factors $\tilde{q}$ and $\hat{q}$ will still measure the Kerr quality of the space-time under analysis. It

may also happen that the space-time is asymptotically flat, in which case we can also use q . In all cases it should be kept in mind, however, that values of q , $\tilde{q}$ and $\hat{q}$ close to 1 will be relevant only in combination with the smallness of these Ricci terms —compared with some characteristic length squared—.

An obvious example to test the previous arguments is the Kerr-de Sitter space-time. Another important example is the rigidly rotating perfect-fluid solution due to Wahlquist, which happens to contain the Kerr spacetime as a particular case. These spacetimes are discussed in the last section where the quality factors are explicitly computed and plotted.

5.3.2 The case without Ernst potential: $\zeta_{[b}R_{a]c}\zeta^c \neq 0$

This is relevant for example for approximate solutions of the Ricci-flat equations, where despite the fact that the Ricci components are assumed to be small one has no extra control on them. It is also relevant for more general situations.

In this case there arise some doubts about the factors κ and $\varkappa$. One possibility to provide them with a reasonable meaning is to also require that the deviations from having an Ernst potential is very small. To that end, recall that in a general space-time the Ernst one-form satisfies [23]

$$\nabla_{[a}\sigma_{b]} = i\eta_{abcd}\zeta^c R^d{}_e \zeta^e$$

which in particular implies that $\nabla_{[a}\sigma_{b]}$ is totally orthogonal to ζ^b . Thus, the quantity $\nabla_{[a}\sigma_{b]}\nabla^{[a}\bar{\sigma}^{b]}$ is non-negative and vanishes if and only if so does the exterior differential of σ_a . We can thus use the following scalar

$$q_\sigma \equiv 1 - \frac{\lambda^2 \nabla_{[a}\sigma_{b]}\nabla^{[a}\bar{\sigma}^{b]}}{\gamma^{ac}\gamma^{bd}\nabla_a\sigma_b\nabla_c\bar{\sigma}_d}$$

as an appropriate quality factor measuring the vanishing of $\nabla_{[a}\sigma_{b]}\nabla^{[a}\bar{\sigma}^{b]}$, and thereby the local existence of an Ernst potential. Observe that $q_\sigma = 1$ if and only if there exists σ such that $\sigma_a = \nabla_a\sigma$ locally. Then, one can use either of the following quality factors for general stationary spacetimes

$$\tilde{q}q_\sigma, \quad \hat{q}q_\sigma$$

keeping in mind that only values of q_σ near one provide a reliable result.

To test the behavior of this refined quality factors we have used the standard Lense-Thirring approximate solution for a rotating stationary isolated body and the Kerr-Newman solution of the Einstein-Maxwell equations, see Section 6.

6 Examples

In this section we present explicit numerical studies of the quality factors introduced above.

6.1 The Curzon-Chazy solution

The Curzon-Chazy solution is a vacuum solution to the Einstein equations which is asymptotically flat. It is the simplest member of the *Weyl class* (see [33] for further details). The explicit form of the metric is

$$ds^2 = -e^{-\frac{2M}{R}}dT^2 + e^{\frac{2M}{R}}\left(e^{-\frac{M^2\sin^2\theta}{R^2}}(dR^2 + R^2d\theta^2) + R^2\sin^2\theta d\phi^2\right), \quad M \in \mathbb{R} \quad (65)$$

The metric functions are everywhere regular except at $R = 0$. The structure of this singularity is rather complicated, being a directional singularity, and it has been studied by a number of authors (see [12] and references therein for a good account of this). In the present study we restrict our attention to the region $-\infty < T < \infty$, $0 < R$, $0 < \theta < \pi$ and $0 < \phi < 2\pi$. In this region there exists an asymptotically flat end $\mathcal{M}_\infty$ located at $\{R > a\}$, with $a \in \mathbb{R}$ big enough and $\partial/\partial T$ as the asymptotic time translation. Therefore we can use the quality factor q




Figure 1: Graphical representation of the quality factor q for the Curzon-Chazy solution when $M = 12$. In this picture we have defined the variables $y = R \cos \theta$, $x = R \sin \theta$ so the vertical line $x = 0$ is the axis of the axial Killing vector $\partial/\partial\phi$. The curves are contour lines with constant q and the contour interval is $1/100$. We can indeed obtain a contour surface by rotating each contour curve around the axial Killing vector axis. The white central region is a set of points where $q < 1/100$; actually, $q = 0$ on the axis for $R \leq M (= 12)$. On the other hand, the outer white region is a set of points where $q > 99/100$; in fact, $q \rightarrow 1$ when $R \rightarrow \infty$. The plot indicates the presence of points whithin the axial Killing vector axis where q is not differentiable. These are probably due to the use of square roots in the denominator of q , nevertheless q is perfectly continuous everywhere.

introduced in Definition 1 to compute the proximity of this solution to the Kerr space-time. In fact the explicit form of q for this solution is

$$q = 1 - \frac{M^2 \sin^2 \theta \left(e^{\frac{2M}{R}} - 1 \right)^2 (M^2 + 3R^2) - 3M^3 R \sin^2 \theta \left(e^{\frac{4M}{R}} - 1 \right) + 3R^2 \left(e^{\frac{2M}{R}} (M - R) + M + R \right)^2}{\left(\left(e^{\frac{2M}{R}} - 1 \right) \sqrt{M^2 \sin^2 \theta (M^2 - 3MR + 3R^2) + 3R^2 (M - R)^2 + 2\sqrt{3}MR} \right)^2},$$

where $\vec{\zeta} = \partial/\partial T$. In figure 1 we present a graph of this quantity for a given value of the parameter M . Observe that q reduces at the axis of symmetry ($\theta = 0$) to

$$q|_{\theta=0} = 1 - \frac{\left(e^{\frac{2M}{R}} (M - R) + M + R \right)^2}{\left(e^{\frac{2M}{R}} |M - R| - |M - R| + 2M \right)^2}$$

so that $q|_{\theta=0} = 0$ at the axis for $R \leq M$, but $q|_{\theta=0} \neq 0$ at the axis for $R > M$. In addition, $q \rightarrow 1$ when $R \rightarrow \infty$.

6.2 The $\delta = 2$ Tomimatsu-Sato solution

The Tomimatsu-Sato space-time [34] can be described by a metric whose non-vanishing components are given by

$$\begin{aligned} g_{tt} &= -\frac{A(x, y)}{B(x, y)}, \quad g_{t\phi} = -\frac{2MqC(x, y)(-1 + y^2)}{B(x, y)}, \quad g_{xx} = \frac{M^2 B(x, y)}{p^2 \delta^2 (-1 + x^2)(x^2 - y^2)^3}, \\ g_{yy} &= \frac{M^2 B(x, y)}{p^2 \delta^2 (-1 + y^2)(-x^2 + y^2)^3}, \\ g_{\phi\phi} &= -\frac{M^2 (-1 + y^2)(p^2 B^2(x, y)(-1 + x^2) + 4q^2 \delta^2 C^2(x, y)(-1 + y^2))}{A(x, y)B(x, y)\delta^2}, \end{aligned} \quad (66)$$

M, p, q, δ being constants, and

$$A(x, y) = \left(p^2 (x^2 - 1)^2 + q^2 (1 - y^2)^2 \right)^2 - 4p^2 q^2 (x^2 - 1)(1 - y^2)(x^2 - y^2)^2, \quad (67)$$

$$B(x, y) = (p^2 x^4 + 2px^3 - 2px + q^2 y^4 - 1)^2 + 4q^2 y^2 (px^3 - pxy^2 - y^2 + 1)^2, \quad (68)$$

$$\begin{aligned} C(x, y) &= p^3 x (1 - x^2) (2(x^4 - 1) + (x^2 + 3)(1 - y^2)) + \\ & p^2 (x^2 - 1) ((x^2 - 1)(1 - y^2) - 4x^2 (x^2 - y^2)) + q^2 (1 - y^2)^3 (px + 1). \end{aligned} \quad (69)$$

This is a vacuum solution of the Einstein equations if $\delta = 2$ and $p^2 + q^2 = 1$. The coordinate ranges are $-\infty < t < \infty$, $-1 < y < 1$, $-\infty < x < -1$, $1 < x < \infty$. In the figure 2 a numerical study of the quality factor q for $\zeta = \partial/\partial t$ is shown.

6.3 The Wahlquist perfect-fluid space-time

A rigidly rotating perfect fluid solution to Einstein field equations was found by Wahlquist in [35], see also [36, 31], which is known to have a vanishing Simon tensor [19, 31] and is the unique stationary and axially symmetric such solution of Petrov type D with a particular linear equation of state [30]. The Wahlquist spacetime actually contains the Kerr-de Sitter space-time as a particular limit case [24]. There are several equivalent but different forms for the Wahlquist solution in the literature, see [25, 29] for a discussion of this, and the best choice for our purposes is the following in local coordinates $\{t, \varphi, y, z\}$

$$ds^2 = -\frac{V - U}{v_1 + v_2} \left(dt - \frac{v_1 V + v_2 U}{V - U} d\varphi \right)^2 + \frac{UV(v_1 + v_2)}{V - U} d\varphi^2 + (v_1 + v_2) \left(\frac{dy^2}{V} + \frac{dz^2}{U} \right)$$

where $v_1(z) = (\cosh(2\beta z) - 1)/(2\beta^2)$, $v_2(y) = (1 - \cos(2\beta y))/(2\beta^2)$, and

$$\begin{aligned} U(z) &= Q_0 - \left(\nu_0 + \frac{\mu_0}{\beta^2} \right) v_1 + \frac{\sinh(2\beta z)}{2\beta} \left(a_1 + z \frac{\mu_0}{\beta^2} \right) \\ V(y) &= Q_0 + \left(\nu_0 + \frac{\mu_0}{\beta^2} \right) v_2 + \frac{\sin(2\beta y)}{2\beta} \left(a_2 - y \frac{\mu_0}{\beta^2} \right) \end{aligned}$$

with $Q_0, a_1, a_2, \nu_0, \mu_0$ and β arbitrary constants. The Kerr-de Sitter-NUT line element is obtained by setting $\beta = 0$, and a_1 is essentially the NUT parameter. Thus, good values of the quality factors are only expected for small β and a_1 . For this solution, the space-time Simon tensor vanishes identically and therefore the quantity q_1 (and hence q_2) is identically 1. Also the Ernst 1-form is closed so there is a local Ernst potential (or equivalently $q_\sigma = 1$). This means that $\hat{q}q_\sigma = \tilde{q}q_\sigma = \kappa\kappa$. We provide a numerical study of $\kappa\kappa$ computed for the Killing vector $\vec{\zeta} = \partial/\partial t$ in figure 3.

6.3.1 The Kerr-de Sitter spacetime

As said above the Kerr-de Sitter spacetime can be regarded as a particular case of the Wahlquist solution just discussed. The Kerr-de Sitter limit is obtained if we set the restrictions [24]

$$Q_0 = A^2, \quad \nu_0 = 1 - A^2 \Lambda/3, \quad a_1 = 0, \quad a_2 = -2M, \quad \mu_0 = -\Lambda, \quad \beta = 0,$$




Figure 2: Representation of the Tomimatsu-Sato global structure for $p = 1/5$. The grey region corresponds to those points in which the axial Killing vector $\partial/\partial\phi$ is space-like, so no causality violations are expected there. The variable X is defined by the condition $X = -1/x$. The lines $X = \pm 1$ correspond to conical singularities and the lines $y = \pm 1$ are rotation axes for the Killing vector $\partial/\partial\phi$. The vertical line $X = 0$ is space-like infinity for two different asymptotically flat regions (these are each characterised by $-1 < X < 0$ and $0 < X < 1$). The two black thick closed curves are the ergosurfaces and the two dots are metric (naked) singularities. We have superimposed some contour curves for the quality factor q indicating the corresponding value of q on each curve. In this particular case this scalar turns out to be positive everywhere (and it approaches $q_\infty = 1$ at $X = 0$).

where A , M and Λ correspond to the angular momentum per unit mass, the mass and the cosmological constant of the Kerr-de Sitter spacetime, respectively. In figure 4 we show a plot of the quality factor $\hat{q} = \tilde{q} = \kappa\kappa$ computed for the Killing vector $\tilde{\zeta} = \partial/\partial t$.

6.4 The Kerr-Newman spacetime

Consider the Kerr-Newman spacetime in the standard Boyer-Linquist coordinates

$$\begin{aligned}
 g_{tt} &= \frac{2MR - A^2 \cos^2 \theta - Q^2 - R^2}{R^2 + A^2 \cos^2 \theta}, & g_{t\phi} &= -\frac{A(2MR - Q^2) \sin^2 \theta}{R^2 + A^2 \cos^2 \theta}, \\
 g_{\theta\theta} &= R^2 + A^2 \cos^2 \theta, & g_{\phi\phi} &= \frac{\sin^2 \theta ((A^2 + R^2)^2 - A^2(A^2 + Q^2 + R^2 - 2MR) \sin^2 \theta)}{R^2 + A^2 \cos^2 \theta}, \\
 g_{RR} &= \frac{R^2 + A^2 \cos^2 \theta}{A^2 + Q^2 + R^2 - 2MR}
 \end{aligned} \tag{70}$$

which is a solution of the Einstein-Maxwell equations. For the Kerr-Newman spacetime the Ernst 1-form is not closed and hence one cannot define the Ernst potential as used here. However, as is well known [33], one can define a complex potential—in general Einstein-Maxwell spacetimes—for a linear combination of the twist one-form ω_c with a one-form associated to the complex




Figure 3: In this figure we show a representation of the quality factor $\hat{q} = \tilde{q} = \kappa\kappa$ for the Wahlquist solution with parameters $Q_0 = 4$, $a_1 = 3.015$, $a_2 = 1.585$, $\nu_0 = 0.64$, $\mu_0 = 1.265$, $\beta = 1.88$ and $\zeta = \partial/\partial t$. Since we are only interested in the region which is close to Kerr, only contour curves at which $\hat{q} > 90/100$ are represented (in this case the contour interval is $5/1000$). The picture on the left is a representation of the factor $\kappa\kappa$ in the plane $z - y$. To facilitate the understanding of this graph, the coordinate z has been compactified by means of the transformation $z \rightarrow \tan z$ and the grey region is the region where ζ is causal. The black vertical lines correspond to points where the determinant spanned by the Killing vector vanishes (note that the leftmost of these lines serves as the boundary of the grey region). The pattern shown in this picture repeats itself *quasi-periodically* along the y -axis. The black dots within the grey region represent points where $\hat{q} > 90/100$. The graph on the right is an inset of a region containing the black dots, made to better appreciate the structure of such region. These numerical representations indicate that $\hat{q}$ reaches the unity value only at isolated points within the studied domain.

self-dual Maxwell 2-form and the Killing vector. Using this generalized potential one can define generalizations of the Mars-Simon tensor and find characterizations of the Kerr-Newman space-time analogous to those used in this paper for the Kerr case: see [37] for the details. Given that the Kerr-Newman space-time is asymptotically flat one can thereby define a quality factor completely analogous to q . This is, however, outside the scope of this paper.

On the other hand, we can pursue our program and compute the quality factors $\tilde{q}q_\sigma$ and $\hat{q}q_\sigma$ for the Kerr-Newman solution. It turns out that $q_1 = q_2 = 1$ as one can explicitly check, and therefore, the quality factor which we need to study is $\tilde{q}q_\sigma = \hat{q}q_\sigma = q_\sigma\kappa\kappa$. In this case one can compute the factors κ and κ in a closed form getting

$$\kappa = 1 - \frac{Q^4}{(A^2 \cos(2\theta) + A^2 + 2R(R - 2M) + 3Q^2)^2}, \quad (71)$$

$$\kappa = \begin{cases} \frac{(A^2 M \cos^2 \theta + R(MR - Q^2))^2}{(A^2 \cos^2 \theta + R^2)(A^2 M^2 \cos^2 \theta + (Q^2 - MR)^2)}, & \text{sign}\left(\mathbf{g}\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right)\right) < 0, \\ \frac{A^2 Q^4 \cos^2 \theta}{(A^2 \cos^2 \theta + R^2)(A^2 M^2 \cos^2 \theta + (Q^2 - MR)^2)}, & \text{sign}\left(\mathbf{g}\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right)\right) > 0. \end{cases} \quad (72)$$

Note that in the limit $Q \rightarrow 0$, κ goes to 1 only outside the ergosphere. This does not contradict the proposition 4 as it is formulated under the assumption $\mathbf{g}(\partial/\partial t, \partial/\partial t) < 0$. In the region $\mathbf{g}(\partial/\partial t, \partial/\partial t) > 0$, κ tends to 0 when Q approaches zero. A numerical study of $q_\sigma\kappa\kappa$ for some selected cases is shown in figure 5.




Figure 4: In these figures we show a representation of the quality factor $\hat{q} = \tilde{q} = \kappa\chi$ for the Kerr-de Sitter solution with parameters $M = 1.5$, $A = 1.32$ and Killing vector $\vec{\zeta} = \partial/\partial t$. In the left picture $\Lambda = 0.1$ and in the right picture $\Lambda = 1$. We have represented the quality factor in the plane $y - z$ by contour curves (contour interval $1/1000$) and only curves with values greater than $99/100$ are represented (this is the region we regard as “close” to unity). The coordinates have been compactified through the relations $y \rightarrow \tan y$, $z \rightarrow \tan z$. In the graphs, $y \in (-\pi/2, \pi/2)$, $z \in (0, \pi/2)$ (the part with $z \in (-\pi/2, 0)$ is obtained by a reflection under the axis $z = 0$). We have also included a representation of the set of points in which the Killing vector $\partial/\partial t$ is timelike (shaded region) and the points where the determinant spanned by the Killing vectors $\partial/\partial t$, $\partial/\partial \phi$ is vanishing (vertical lines and horizontal line). There is a region where the quality factor is above $999/1000$ which is bounded by the thick contour line (it contains the singularity $y = 0, z = 0$ as an accumulation point). This region is larger in the left picture as Λ is smaller in this case. One can check that the size of this region increases when Λ approaches zero.

6.5 The Lense-Thirring approximation

The Lense-Thirring metric can be regarded as a first order approximation to the (exterior) metric of a rotating compact body. The metric tensor components are obtained from (21) by just retaining the leading terms

$$g_{tt} = -1 + \frac{2M}{r}, \quad g_{tx^i} = -\epsilon_{ijk} \frac{4S^i x^k}{r^3}, \quad g_{x^i x^j} = \delta_{ij}. \quad (73)$$

Without loss of generality we may adopt the choice $S^1 = S^2 = 0$ (i.e. we choose a coordinate system adapted to the body rotation axis). For our computations, it is convenient to express the above metric in spherical coordinates (R, θ, ϕ) which are related to the coordinates (x^1, x^2, x^3) through the usual relation $x^1 = R \sin \theta \cos \phi$, $x^2 = R \sin \theta \sin \phi$, $x^3 = R \cos \theta$. We present in figure 6 a numerical study of the quality factor $q_\sigma \hat{q}$ for a certain choice of M and of the non-vanishing component S^3 .

7 Conclusions

Taking the invariant characterizations of the Kerr solution written in terms of the Mars-Simon and the space-time Simon tensors we have constructed a number of dimensionless, non-negative scalar quantities (quality factors) which attain the value $+1$ in a Ricci flat space-time if and only if it is locally isometric to the Kerr solution. These quality factors depend on the existence of a time-like Killing vector in the space-time and as such they can only be used on stationary




Figure 5: Contour curves for the quality factor $\tilde{q}q_\sigma = \hat{q}q_\sigma = q_\sigma\kappa\kappa$ for the Kerr Newman solution with $\vec{\zeta} = \partial/\partial t$. The coordinate R (horizontal axis) has been compactified with the transformation $R \rightarrow \tan R$ and the coordinate θ runs along the vertical axis (hence the upper and lower parts of the plots represent the axis of the Killing field $\partial/\partial\phi$ and they should be identified). The values of the mass and angular momentum per unit mass are $M = 0.4$, $A = 0.35$ and we consider two different values of the charge parameter Q . On the graph of the left we set $Q = 0.19$ and in this case there are two Killing horizons represented by the two parallel vertical lines. The dashed curves are the ergosurfaces and the dot is the ring singularity located at $R = 0$, $\theta = 0$. Only contour curves with values of the quality factor greater than $99/100$ have been represented (the contour interval is $1/10000$). On the graph of the right $Q = 4$ and there are neither Killing horizons, nor ergosurfaces, but the central dot representing the ring singularity remains. In this case the contour interval is $1/100$ and no further restrictions on the contour curves have been placed. In both cases the quality factor approaches unity for $R \rightarrow \pm\pi/2$ (the asymptotically flat regions).

space-times. However, we must stress that in all the examples studied in section 6 the positivity properties of the quality factors were kept even in those points where the Killing vector is not time-like. This strongly suggests that all the quality factors studied in this paper are indeed non-negative regardless of the causal character of the Killing vector entering in their definition. A rigorous proof of this fact has eluded us so far.

Another interesting question regarding the quality factors is whether they fulfill some hyperbolic differential equation similar to eq. (19) for the Mars-Simon tensor. If that was the case then one could analyse the Cauchy problem of such an equation and the global existence of its solutions. This could serve to address the non-linear stability of the Kerr solution under the assumption of the existence of a Killing vector.

We have already mentioned in the introduction the uniqueness results dealing with the Kerr solution and the recent progress in this direction. In [1] a uniqueness result for the Kerr solution was proven which applies to smooth asymptotically flat stationary vacuum solutions of the Einstein equations. Besides the usual assumptions of the space-time being the exterior region of a regular black hole, a hypothesis about the space-time being “close” to Kerr is assumed. This “Perturbation of Kerr assumption” states that the components of the Mars-Simon tensor with respect to an orthonormal frame are bounded by a certain quantity when evaluated on a partial Cauchy hypersurface of the *domain of outer communication*. One could entertain the possibility of expressing this assumption in terms of one of the quality factors introduced in this article. This might result in alternative, maybe more powerful, uniqueness results for the Kerr solution.

Finally, there is the open question of how to generalize the quality factors to the generic




Figure 6: The graph of the left is a plot of the contour curves for the quality factor $q_\sigma \hat{q}$ for the case of the approximate Lense-Thirring solution with parameters $M = 2$, $S^1 = S^2 = 0$, $S^3 = 14$ and fixed $\theta = \pi/4$ (t is suppressed so only the $R - \phi$ dependency is kept). As in some of the previous plots, the contour interval is $1/100$. Note that the curves are all vertical lines and one can check that this behaviour is maintained for different values of θ , suggesting that the contour hypersurfaces have axial symmetry, as expected on intuitive grounds. We represent one of these contour hypersurfaces in the graph on the right. In this graph the coordinates x^2 and t have been suppressed and the value of $q_\sigma \hat{q}$ is $7/10$. Due to the axial symmetry one just has to rotate the curve drawn in the graph around the vertical line to obtain the corresponding contour hypersurface. Note the presence of a region in which numerical value of $q_\sigma \hat{q}$ is less than $1/100$ (the inner boundary of this region is $R = 2M = 4$).

situation without Killing vector fields.

8 Acknowledgements

JMMS is supported by grants FIS2010-15492 (MICINN), GIU06/37 (UPV/EHU), P09-FQM-4496 (J. Andalucía—FEDER) and UFI 11/55 (UPV/EHU). AGP is supported by the Research Centre of Mathematics of the University of Minho (Portugal) through the “Fundação para a Ciência e a Tecnologia (FCT) Pluriannual Funding Program” and through project CERN/FP/123609/2011.

References

- [1] S. Alexakis, A. D. Ionescu, and Klainerman S., *Uniqueness of Smooth Stationary Black Holes in Vacuum: Small Perturbations of the Kerr Spaces*, Comm. Math. Phys. **299** (2010), no. 1, 89–127.
- [2] R. Beig and W. Simon, *The Stationary Gravitational Field near Spatial Infinity*, Gen. Rel. Grav. **12** (1980), no. 12, 1003–1013.
- [3] L. Bel, *Sur la radiation gravitationnelle*, C. R. Acad. Sci. Paris **247** (1958), 1094–1096.
- [4] D. Christodoulou and S. Klainerman, *The global nonlinear stability of the Minkowski space*, Princeton Mathematical Series, vol. 41, Princeton University Press, Princeton, NJ, 1993.

- [5] P. T. Chruściel and J. L. Costa, *On uniqueness of stationary vacuum black holes*, Astérisque (2008), no. 321, 195–265, Géométrie différentielle, physique mathématique, mathématiques et société. I.
- [6] P. T. Chruściel, J. Lopes Costa, and M. Heusler, *Stationary black holes: Uniqueness and beyond*, Living Reviews in Relativity **15** (2012), no. 7.
- [7] J. J. Ferrando and J. A. Sáez, *An intrinsic characterization of the Kerr metric*, Class. Quantum Grav. **26** (2009), 075013,13.
- [8] H. Friedrich, *Existence and structure of past asymptotically simple solutions of Einstein’s field equations with positive cosmological constant*, J. Geom. Phys. **3** (1986), no. 1, 101–117.
- [9] ———, *On the existence of n -geodesically complete or future complete solutions of Einstein’s field equations with smooth asymptotic structure*, Comm. Math. Phys. **107** (1986), no. 4, 587–609.
- [10] ———, *On the global existence and the asymptotic behavior of solutions to the Einstein-Maxwell-Yang-Mills equations*, J. Differential Geom. **34** (1991), no. 2, 275–345.
- [11] A. García-Parrado and J. A. Valiente Kroon, *Kerr initial data*, Classical Quantum Gravity **25** (2008), no. 20, 205018, 20.
- [12] J. B. Griffiths and J. Podolský, *Exact space-times in Einstein’s general relativity*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, 2009.
- [13] Markus Heusler, *Black hole uniqueness theorems*, Cambridge Lecture Notes in Physics, vol. 6, Cambridge University Press, Cambridge, 1996.
- [14] A. D. Ionescu and S. Klainerman, *On the uniqueness of smooth, stationary black holes in vacuum*, Invent. Math. **175** (2009), no. 1, 35–102.
- [15] A.D. Ionescu and S. Klainerman, *On the local extension of killing vector-fields in ricci flat manifolds*, <http://arxiv.org/abs/1108.3575> (2011).
- [16] R. P. Kerr, *Gravitational field of a spinning mass as an example of an algebraically special metric*, Phys. Rev. Lett. **11** (1963), no. 5, 237–238.
- [17] S. Klainerman and F. Nicolò, *The evolution problem in general relativity*, Progress in Mathematical Physics, vol. 25, Birkhäuser Boston Inc., Boston, MA, 2003.
- [18] A. Komar, *Covariant conservation laws in General Relativity*, Phys. Rev. **113** (1959) 934–936.
- [19] D. Kramer, *Rigidly rotating perfect fluids*, Astronom. Nachr. **307** (1986), no. 5, 309–312.
- [20] H. Lindblad and I. Rodnianski, *Global existence for the Einstein vacuum equations in wave coordinates*, Comm. Math. Phys. **256** (2005), no. 1, 43–110.
- [21] M. Mars, *A spacetime characterization of the Kerr metric*, Classical Quantum Gravity **16** (1999), no. 7, 2507–2523.
- [22] ———, *Uniqueness properties of the Kerr metric*, Classical Quantum Gravity **17** (2000), no. 16, 3353–3373.
- [23] ———, *Spacetime Ehlers group: transformation law for the Weyl tensor*, Classical Quantum Gravity **18** (2001), no. 4, 719–738.
- [24] ———, *Wahlquist-Newman solution*, Phys. Rev. D (3) **63** (2001), no. 6, 064022, 9.
- [25] M. Mars and J. M. M. Senovilla, *Comment on: “A generalization of the Wahlquist solution” [Internat. J. Modern Phys. D **7** (1998), no. 6, 927–941 by T. Papakostas]*, Internat. J. Modern Phys. D **8** (1999), no. 6, 753–754.

- [26] J. M. Martín-García, *xAct: efficient tensor computer algebra*, <http://www.xact.es>.
- [27] ———, *xPerm: fast index canonicalization for tensor computer algebra*, Computer Physics Communications **179** (2008), 597–603.
- [28] R. Penrose and W. Rindler, *Spinors and space-time. Vol. 1*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, 1987.
- [29] I. Rácz and J. Zsigrai, *Generating new perfect-fluid solutions from known ones*, Classical Quantum Gravity **13** (1996), no. 10, 2783–2795.
- [30] J. M. M. Senovilla, *Stationary axisymmetric perfect-fluid metrics with $q + 3p = \text{const}$* , Phys. Lett. A **123** (1987), no. 5, 211–214.
- [31] ———, *Stationary and axisymmetric perfect-fluid solutions to Einstein's equations*, Rotating objects and relativistic physics (El Escorial, 1992), Lecture Notes in Phys., vol. 423, Springer, Berlin, 1993, pp. 73–88.
- [32] ———, *Super-energy tensors*, Class. Quantum Grav. **17** (2000), no. 14, 2799–2841.
- [33] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers, and E. Herlt, *Exact solutions of Einstein's field equations*, second ed., Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, 2003.
- [34] A. Tomimatsu and H. Sato, *New Exact Solution for the Gravitational Field of a Spinning mass*, Phys. Rev. Lett. **29** (1972), no. 19, 1344–1345.
- [35] H. D. Wahlquist, *Interior Solution for a Finite Rotating Body of Perfect Fluid*, Phys. Rev. **172** (1968), no. 5, 1291–1296.
- [36] ———, *The dyadic approach to solutions for rotating rigid bodies*, Rotating objects and relativistic physics (El Escorial, 1992), Lecture Notes in Phys., vol. 423, Springer, Berlin, 1993, pp. 55–72.
- [37] Willie Wai-Yeung Wong, *A space-time characterization of the Kerr-Newman metric*, Annales Henri Poincaré **10** (2009) 453–484.